



EXHIBIT C
CLAIMS AS PENDING FOLLOWING ENTRY OF
AMENDMENT OF AUGUST 19, 2002
U.S. PATENT APPLICATION NO. 09/823,307
ATTORNEY DOCKET NO. 7853-235-999

RECEIVED
AUG 26 2002
TECH CENTER 1600/2900

21. (Amended) A method of inhibiting costimulation of human T lymphocytes comprising: contacting a human T lymphocyte with a monoclonal antibody that recognizes a human 8F4 polypeptide, wherein said 8F4 polypeptide:

- a) is an inducible T cell costimulatory molecule;
- b) occurs on two-signal-activated human T lymphocytes;
- c) exhibits a molecular weight of about 55 to 60 kilodaltons as determined by non-reducing sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE); and
- d) is a dimer of two peptide chains exhibiting molecular weights of about 27 kilodaltons and 29 kilodaltons, as measured by reducing SDS-PAGE, such that costimulation of the human T lymphocyte is modulated.

25. (Amended) The method of claim 21, wherein the monoclonal antibody recognizes the human 8F4 polypeptide of about 55 kilodaltons to 60 kilodaltons, as determined by non-reducing SDS-PAGE.

26. (Amended) The method of claim 21, wherein the monoclonal antibody recognizes the peptide chain of about 27 kilodaltons, as determined by reducing SDS-PAGE.

27. (Amended) The method of claim 21, wherein the monoclonal antibody recognizes the peptide chain of about 29 kilodaltons, as determined by reducing SDS-PAGE.

28. (Amended) The method of claim 21, wherein the monoclonal antibody recognizes a human 8F4 polypeptide present on activated human CD4⁺ T lymphocytes and activated human CD8⁺ T lymphocytes.

30. (Amended) A method of inhibiting rejection of an organ transplant, comprising: administering to an individual in need of such inhibition an 8F4 inhibitory molecule, which 8F4 inhibitory molecule is a monoclonal antibody that recognizes a human 8F4 polypeptide, wherein said 8F4 polypeptide:

- a) is an inducible T cell costimulatory molecule;
- b) occurs on two-signal-activated human T lymphocytes;
- c) exhibits a molecular weight of about 55 to 60 kilodaltons as determined by non-reducing sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE); and
- d) is a dimer of two peptide chains exhibiting molecular weights of about 27 kilodaltons and 29 kilodaltons, as measured by reducing SDS-PAGE, in an amount sufficient to inhibit rejection of an organ transplant.

32. (Amended) The method of claim 30, wherein the monoclonal antibody recognizes the human 8F4 polypeptide of about 55 kilodaltons to 60 kilodaltons, as determined by non-reducing SDS-PAGE.

33. (Amended) The method of claim 30, wherein the monoclonal antibody recognizes the peptide chain of about 27 kilodaltons, as determined by reducing SDS-PAGE.

34. (Amended) The method of claim 30, wherein the monoclonal antibody recognizes the peptide chain of about 29 kilodaltons, as determined by reducing SDS-PAGE.

35. (Amended) The method of claim 30, wherein the monoclonal antibody recognizes a human 8F4 polypeptide present on activated human CD4⁺ T lymphocytes and activated human CD8⁺ T lymphocytes.